# 3.3.2 Tests for specified microorganisms

This text is based on the internationally-harmonized texts developed by the Pharmacopoeial Discussion Group (PDG). Some editorial modifications have been made in order to be in line with the style used in The International Pharmacopoeia.

The tests described hereafter will allow determination of the absence of, or limited occurrence of, specified microorganisms which may be detected under the conditions described.

The tests are designed primarily to determine whether a substance or preparation complies with an established specification for microbiological quality. When used for such purposes follow the instructions given below, including the number of samples to be taken and interpret the results as stated below.

Alternative microbiological procedures, including automated methods may be used, provided that their equivalence to the pharmacopoeial method has been demonstrated.

## GENERAL PROCEDURES

The preparation of samples is carried out as described in <u>3.3.1 Microbial enumeration tests</u>.

If the product to be examined has antimicrobial activity, this is insofar as possible removed or neutralized as described in <u>3.3.1</u> <u>Microbial enumeration tests</u>.

If surface-active substances are used for sample preparation, their absence of toxicity for microorganisms and their compatibility with inactivators used must be demonstrated as described in <u>3.3.1 Microbial enumeration tests</u>.

# GROWTH PROMOTING AND INHIBITORY PROPERTIES OF THE MEDIA, SUITABILITY OF THE TEST AND NEGATIVE CONTROLS

The ability of the test to detect microorganisms in the presence of the product to be tested must be established. Suitability must be confirmed if a change in testing performance, or the product, which may affect the outcome of the test is introduced.

#### Preparation of test strains

Use standardized stable suspensions of test strains or prepare as stated below. Seed-lot culture maintenance techniques (seed-lot systems) are used so that the viable microorganisms used for inoculation are not more than 5 passages removed from the original master seed-lot.

#### Aerobic microorganisms

Grow each of the bacterial test strains separately in containers containing *casein soya bean digest broth* or on *casein soya bean digest agar* at 30–35 °C for 18–24 h. Grow the test strain for *Candida albicans* separately on *Sabouraud-dextrose agar* or in *Sabouraud-dextrose broth* at 20–25 °C for 2–3 days.

Staphylococcus aureus such as ATCC 6538, NCIMB 9518, CIP 4.83 or NBRC 13276,

Pseudomonas aeruginosa such as ATCC 9027, NCIMB 8626, CIP 82.118 or NBRC 13275,

Escherichia coli such as ATCC 8739, NCIMB 8545, CIP 53.126 or NBRC 3972,

Salmonella enterica subsp. enterica serovar Typhimurium such as ATCC 14028 or, as an alternative, Salmonella enterica subsp. enterica serovar Abony such as NBRC 100797, NCTC 6017 or CIP 80.39,

Candida albicans such as ATCC 10231, NCPF 3179, IP 48.72 or NBRC 1594.

Use *buffered sodium chloride-peptone solution* pH 7.0 or *phosphate buffer pH 7.2* to make test suspensions. Use the suspensions within 2 h or within 24 h if stored at 2–8 °C.

#### Clostridia

Use *Clostridium sporogenes* such as ATCC 11437 (NBRC 14293, NCIMB 12343, CIP 100651) or ATCC 19404 (NCTC 532 or CIP 79.3). Grow the clostridial test strain under anaerobic conditions in *reinforced medium for Clostridia* at 30–35 °C for 24–48 h. As an alternative to preparing and then diluting down a fresh suspension of vegetative cells of *CI. sporogenes* a stable spore suspension is used for test inoculation. The stable spore suspension may be maintained at 2–8 °C for a validated period.

# Negative control

To verify testing conditions a negative control is performed using the chosen diluent in place of the test preparation. There must be no growth of microorganisms. A negative control is also performed when testing the products as described under *Testing of products*. A failed negative control requires an investigation.

#### Growth promotion and inhibitory properties of the media

Test each batch of ready-prepared medium and each batch of medium prepared either from dehydrated medium or from ingredients.

Verify suitable properties of relevant media as described in Table 1.

Medium	Property	Test strains		
Test for bile-tolerant gram-negative bacteria				
Enterobacteria enrichment broth-Mossel	Growth-promoting	E. coli P. aeruginosa		
	Inhibitory	S. aureus		
Violet red bile glucose agar	Growth-promoting + Indicative	E. coli P. aeruginosa		
Test for Escherichia coli				
MacConkov broth	Growth-promoting	E. coli		
	Inhibitory	S. aureus		
MacConkey agar	Growth-promoting + Indicative	E. coli		
Test for Salmonella				
Rappaport Vassiliadis Salmonella enrichment broth	Growth-promoting	Salmonella enterica subsp. enterica serovar Typhimurium or Salmonella enterica subsp. enterica serovar Abony		
	Inhibitory	S. aureus		
Xylose, lysine, deoxycholate agar	Growth-promoting + Indicative	Salmonella enterica subsp. enterica serovar Typhimurium or Salmonella enterica subsp. enterica serovar Abony		
Test for Pseudomonas aeruginosa				
Cetrimide agar	Growth-promoting	P. aeruginosa		
	Inhibitory	E. coli		
Test for Staphylococcus aureus				
Mannitol salt agar	Growth-promoting + Indicative	S. aureus		
	Inhibitory	E. coli		
Test for <i>Clostridia</i>				
Reinforced medium for Clostridia	Growth-promoting	Cl. sporogenes		
Columbia agar	Growth-promoting	Cl. sporogenes		
Test for Candida albicans				
Sabouraud dextrose broth	Growth-promoting	C. albicans		
Sabouraud dextrose broth	Growth-promoting + Indicative	C. albicans		

*Test for growth-promoting properties, liquid media.* Inoculate a portion of the appropriate medium with a small number (not more than 100 CFU) of the appropriate microorganism. Incubate at the specified temperature for not more than the shortest period of time specified in the test. Clearly visible growth of the microorganism comparable to that previously obtained with a previously tested and approved batch of medium occurs.

*Test for growth-promoting properties, solid media.* Perform surface-spread method, inoculating each plate with a small number (not more than 100 CFU) of the appropriate microorganism. Incubate at the specified temperature for not more than the shortest period of time specified in the test. Growth of the microorganism comparable to that previously obtained with a previously tested and approved batch of medium occurs.

*Test for inhibitory properties, liquid or solid media.* Inoculate the appropriate medium with at least 100 CFU of the appropriate microorganism. Incubate at the specified temperature for not less than the longest period of time specified in the test. No growth of the test microorganism occurs.

*Test for indicative properties.* Perform surface-spread method, inoculating each plate with a small number (not more than 100 CFU) of the appropriate microorganism. Incubate at the specified temperature for a period of time within the range specified in

the test. Colonies are comparable in appearance and indication reactions to those previously obtained with a previously tested and approved batch of medium.

## Suitability of the test method

For each product to be tested perform a sample preparation as described in the relevant paragraph in the section below on *Testing of products*. Add each test strain at the time of mixing, in the prescribed growth medium. Inoculate the test strains individually. Use a number of microorganisms equivalent to not more than 100 CFU in the inoculated test preparation.

Perform the test as described in the relevant paragraph in *Testing of products* section using the shortest incubation period prescribed.

The specified microorganisms must be detected with the indication reactions as described below under *Testing of products*.

Any antimicrobial activity of the product necessitates a modification of the test procedure (see Neutralization/removal of antimicrobial activity of <u>3.3.1 Microbial enumeration tests</u>).

If for a given product the antimicrobial activity with respect to a microorganism for which testing is prescribed cannot be neutralized, then it is to be assumed that the inhibited microorganism will not be present in the product.

## **TESTING OF PRODUCTS**

#### Bile-tolerant gram-negative bacteria

#### Sample preparation and pre-incubation

Prepare a sample using a 1 in 10 dilution of not less than 1 g of the product to be examined as described in <u>3.3.1 Microbial</u> <u>enumeration tests</u>, but using *casein soya bean digest broth* as the chosen diluent, mix and incubate at 20–25 °C for a time sufficient to resuscitate the bacteria but not sufficient to encourage multiplication of the organisms (usually 2 h but not more than 5 h).

## Test for absence

Unless otherwise prescribed use the volume corresponding to 1g of the product, as prepared in *Sample preparation and pre-incubation*, to inoculate *enterobacteria enrichment broth-Mossel*. Incubate at 30–35 °C for 24–48 h. Subculture on plates of *violet red bile glucose agar*. Incubate at 30–35 °C for 18–24 h.

The product complies with the test if there is no growth of colonies.

#### Quantitative test

Selection and subculture. Inoculate suitable quantities of *enterobacteria enrichment broth-Mossel* with the preparation as described under *Sample preparation and pre-incubation* and/or dilutions of it containing, respectively 0.1 g, 0.01 g and 0.001 g (or 0.1 mL, 0.01 mL and 0.001 mL) of the product to be examined. Incubate at 30–35 °C for 24–48 h. Subculture each of the cultures on a plate of *violet red bile glucose agar*. Incubate at 30–35 °C for 18–24 h.

*Interpretation.* Growth of colonies constitutes a positive result. Note the smallest quantity of the product that gives a positive result and the largest quantity that gives a negative result. Determine from Table 2 the probable number of bacteria.

#### Table 2. Interpretation of results

Results for each quantity of product		of product	Probable number of bacteria per gram or mL of product
0.1 g or 0.1 mL	0.01 g or 0.01 mL	0.001 g or 0.001 mL	
+	+	+	more than 10 <sup>3</sup>
+	+	-	less than 10 <sup>3</sup> and more than 10 <sup>2</sup>
+	-	-	less than 10 <sup>2</sup> and more than 10
-	-	-	less than 10

#### Escherichia coli

Sample preparation and pre-incubation

Prepare a sample using a 1 in 10 dilution of not less than 1 g of the product to be examined as described in <u>3.3.1 Microbial</u> <u>enumeration tests</u> and use 10 mL or the quantity corresponding to 1 g or1 mL to inoculate a suitable amount (determined as described under *Suitability of the test method*) of *casein soya bean digest broth*, mix and incubate at 30–35 °C for 18–24 h.

## Selection and subculture

Shake the container, transfer 1 mL of *casein soya bean digest broth* to 100 mL of *MacConkey broth* and incubate at 42–44 °C for 24–48 h. Subculture on a plate of *MacConkey agar* at 30–35 °C for 18–72 h.

## Interpretation

Growth of colonies indicates the possible presence of E. coli. This is confirmed by identification tests.

The product complies with the test if no colonies are present or if the identification tests are negative.

## Salmonella

## Sample preparation and pre-incubation

Prepare the product to be examined as described in <u>3.3.1 Microbial enumeration tests</u> and use the quantity corresponding to not less than 10 g or 10 mL to inoculate a suitable amount (determined as described under *Suitability of the test method*) of *casein soya bean digest broth*, mix and incubate at 30–35 °C for 18–24 h.

## Selection and subculture

Transfer 0.1 mL of *casein soya bean digest broth* to 10 mL of *Rappaport Vassiliadis Salmonella enrichment broth* and incubate at 30–35 °C for 18–24 h. Subculture on plates of *xylose, lysine, deoxycholate agar.* Incubate at 30–35 °C for 18–48 h.

#### Interpretation

The possible presence of *Salmonella* is indicated by the growth of well-developed, red colonies, with or without black centres. This is confirmed by identification tests.

The product complies with the test if colonies of the types described are not present or if the confirmatory identification tests are negative.

## Pseudomonas aeruginosa

## Sample preparation and pre-incubation

Prepare a sample using a 1 in 10 dilution of not less than 1 g of the product to be examined as described in <u>3.3.1 Microbial</u> <u>enumeration tests</u> and use 10 mL or the quantity corresponding to 1 g or 1 mL to inoculate a suitable amount (determined as described under *Suitability of the test method*) of *casein soya bean digest broth* and mix. When testing transdermal patches filter the volume of sample corresponding to 1 patch of the preparation described under Preparation of the sample in <u>3.3.1 Microbial</u> <u>enumeration tests</u> through a sterile filter membrane and place in 100 mL of *casein soya bean digest broth*. Incubate at 30–35 °C for 18–24 h.

#### Selection and subculture

Subculture on a plate of *cetrimide agar* and incubate at 30–35 °C for 18–72 h.

#### Interpretation

Growth of colonies indicates the possible presence of *P. aeruginosa*. This is confirmed by identification tests.

The product complies with the test if colonies are not present or if the confirmatory identification tests are negative.

#### Staphylococcus aureus

#### Sample preparation and pre-incubation

Prepare a sample using a 1 in 10 dilution of not less than 1 g of the product to be examined as described in <u>3.3.1 Microbial</u> <u>enumeration tests</u> and use 10 mL or the quantity corresponding to 1 g or 1 mL to inoculate a suitable amount (determined as described under *Suitability of the test method*) of *casein soya bean digest broth* and homogenize. When testing transdermal patches filter the volume of sample corresponding to 1 patch of the preparation described under Preparation of the sample in 3.3.1 Microbial enumeration teststhrough a sterile filter membrane and place in 100 mL of *casein soya bean digest broth*. Incubate at 30–35 °C for 18–24 h.

#### Selection and subculture

Subculture on a plate of mannitol salt agar and incubate at 30–35 °C for 18–72 h.

#### Interpretation

The possible presence of S. aureus is indicated by the growth of yellow/white colonies surrounded by a yellow zone. This is

## confirmed by identification tests.

The product complies with the test if colonies of the types described are not present or if the confirmatory identification tests are negative.

## Clostridia

## Sample preparation and heat treatment

Prepare a sample using a 1 in 10 dilution (with a minimum total volume of 20 mL) of not less than 2 g or 2 mL of the product to be examined as described in <u>3.3.1 Microbial enumeration tests</u>. Divide the sample into two portions of at least 10 mL. Heat 1 portion at 80 °C for 10 minutes and cool rapidly. Do not heat the other portion.

## Selection and subculture

Use 10 mL or the quantity corresponding to 1 g or 1 mL of the product to be examined of both portions to inoculate suitable amounts (determined as described under *Suitability of the test method*) of *Reinforced clostridium medium*. Incubate under anaerobic conditions at 30–35 °C for 48 h. After incubation, make subcultures from each container on *Columbiaagar* and incubate under anaerobic conditions at 30–35 °C for 48–72 h.

## Interpretation

The occurrence of anaerobic growth of rods (with or without endospores) giving a negative catalase reaction indicates the presence of *Clostridia*. This is confirmed by identification tests.

The product complies with the test if colonies of the types described are not present or if the confirmatory identification tests are negative.

## Candida albicans

# Sample preparation and pre-incubation

Prepare the product to be examined as described in <u>3.3.1 Microbial enumeration tests</u> and use 10 mL or the quantity corresponding to not less than 1 g or 1 mL to inoculate 100 mL of *Sabouraud-dextrose broth* and mix. Incubate at 30–35 °C for 3–5 days.

#### Selection and subculture

Subculture on a plate of Sabouraud-dextrose agar and incubate at 30-35 °C for 24-48h.

Interpretation

Growth of white colonies may indicate the presence of C. albicans. This is confirmed by identification tests.

The product complies with the test if such colonies are not present or if the confirmatory identification tests are negative.

# RECOMMENDED TEST SOLUTIONS AND CULTURE MEDIA

The following test solutions and culture media have been found satisfactory for the purposes for which they are prescribed in the test for microbial contamination in the pharmacopoeia. Other media may be used provided that their suitability can be demonstrated.

Stock buffer solution. Transfer 34 g of potassium dihydrogen phosphate to a 1000 mL volumetric flask, dissolve in 500 mL of purified water, adjust to pH 7.2  $\pm$  0.2 with sodium hydroxide, add purified water to volume and mix. Dispense in containers and sterilize. Store at a temperature of 2–8 °C.

Phosphate buffer pH 7.2. Prepare a mixture of purified water and stock buffer solution (800:1 V/V) and sterilize.

-Buffered sodium chloride-peptone solution pH 7.0

Potassium dihydrogen phosphate 3.6 g

Disodium hydrogen phosphate dihydrate 7.2 g equivalent to 0.067 M phosphate

Sodium chloride 4.3 g

Peptone (meat or casein) 1.0 g

Purified water 1000 mL

Sterilize in an autoclave using a validated cycle.

-Casein soya bean digest broth

Pancreatic digest of casein 17.0 g

Papaic digest of soya bean 3.0 g

Sodium chloride 5.0 g

Dipotassium hydrogen phosphate 2.5 g

Glucose monohydrate 2.5 g

Purified water 1000 mL

Adjust the pH so that after sterilization it is 7.3 ± 0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

-Casein soya bean digest agar

Pancreatic digest of casein 15.0 g

Papaic digest of soya bean 5.0 g

Sodium chloride 5.0 g

Agar 15.0 g

Purified water 1000 mL

Adjust the pH so that after sterilization it is 7.3 ± 0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

-Sabouraud-dextrose agar

Dextrose 40.0 g

Mixture of peptic digest of animal tissue and

pancreatic digest of casein (1:1) 10.0 g

Agar 15.0 g

Purified water 1000 mL

Adjust the pH so that after sterilization it is 5.6 ± 0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

# -Potato dextrose agar

Infusion from potatoes 200 g

Dextrose 20.0 g

Agar 15.0 g

Purified water 1000 mL

Adjust the pH so that after sterilization it is 5.6 ± 0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

-Sabouraud-dextrose broth

Dextrose 20.0 g

Mixture of peptic digest of animal tissue and

pancreatic digest of casein (1:1) 10.0 g

Purified water 1000 mL

Adjust the pH so that after sterilization it is 5.6  $\pm$  0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

-Enterobacteria enrichment broth-Mossel

Pancreatic digest of gelatin 10.0 g

Glucose monohydrate 5.0 g

Dehydrated ox bile 20.0 g

Potassium dihydrogen phosphate 2.0 g

Disodium hydrogen phosphate dihydrate 8.0 g

Brilliant green 15 mg

Purified water 1000 mL

Adjust the pH so that after heating it is 7.2  $\pm$  0.2 at 25 °C. Heat at 100 °C for 30 minutes and cool immediately.

-Violet red bile glucose agar

Yeast extract 3.0 g

Pancreatic digest of gelatin 7.0 g

Bile salts 1.5 g

Sodium chloride 5.0 g

Glucose monohydrate 10.0 gl

Agar 15.0 g

Neutral red 30 mg

Crystal violet 2 mg

Purified water 1000 mL

Adjust the pH so that after heating it is 7.4 ± 0.2 at 25 °C. Heat to boiling; do not heat in an autoclave.

#### -MacConkey broth

Pancreatic digest of gelatin 20.0 g

Lactose monohydrate 10.0 g

Dehydrated ox bile 5.0 g

Bromocresol purple 10 mg

Purified water 1000 mL

Adjust the pH so that after sterilization it is 7.3 ± 0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

#### -MacConkey agar

Pancreatic digest of gelatin 17.0 g

Peptones (meat and casein) 3.0 g

Lactose monohydrate 10.0 g

Sodium chloride 5.0 g

Bile salts 1.5 g

Agar 13.5 g

Neutral red 30.0 mg

Crystal violet 1 mg

Purified water 1000 mL

Adjust the pH so that after sterilization it is 7.1  $\pm$  0.2 at 25 °C. Boil for 1 minute with constant shaking then sterilize in an autoclave using a validated cycle.

-Rappaport Vassiliadis Salmonella Enrichment Broth

Soya peptone 4.5 g

Magnesium chloride hexahydrate 29.0 g

Sodium chloride 8.0 g

Dipotassium phosphate 0.4 g

Potassium dihydrogen phosphate 0.6 g

Malachite green 0.036 g

Purified water 1000 mL

Dissolve, warming slightly. Sterilize in an autoclave using a validated cycle, at a temperature not exceeding 115 °C. The pH is to be  $5.2 \pm 0.2$  at 25 °C after heating and autoclaving.

-Xylose, lysine, deoxycholate agar

Xylose 3.5 g

L-Lysine 5.0 g

Lactose monohydrate 7.5 g

Sucrose 7.5 g

Sodium chloride 5.0 g

Yeast extract 3.0 g

Phenol red 80 mg

Agar 13.5 g

Sodium deoxycholate 2.5 g

Sodium thiosulfite 6.8 g

Ferric ammonium citrate 0.8 g

Purified water 1000 mL

Adjust the pH so that after heating it is 7.4  $\pm$  0.2 at 25 °C. Heat to boiling, cool to 50 °C and pour into Petri dishes. Do not heat in an autoclave.

#### -Cetrimide agar

Pancreatic digest of gelatin 20.0 g

Magnesium chloride 1.4 g

Dipotassium sulfate 10.0 g

Cetrimide 0.3 g

Agar 13.6 g

Purified water 1 000 mL

Glycerol 10.0 mL

Heat to boiling for 1 minute with shaking. Adjust the pH so that after sterilization it is  $7.2 \pm 0.2$  at 25 °C. Sterilize in an autoclave using a validated cycle.

-Mannitol salt agar

Pancreatic digest of casein 5.0 g Peptic digest of animal tissue 5.0 g Beef extract 1.0 g D-Mannitol 10.0 g Sodium chloride 75.0 g Agar 15.0 g

# Phenol red 0.025 g

Purified water 1000 mL

Heat to boiling for 1 minute with shaking. Adjust the pH so that after sterilization it is 7.4  $\pm$  0.2 at 25 °C. Sterilize in an autoclave using a validated cycle.

-Reinforced medium for Clostridia

Beef extract 10.0 g

Peptone 10.0 g

Yeast extract 3.0 g

Soluble starch 1.0 g

Glucose monohydrate 5.0 g

Cysteine hydrochloride 0.5 g

Sodium chloride 5.0 g

Sodium acetate 3.0 g

Agar 0.5 g

Purified water 1000 mL

Hydrate the agar, dissolve by heating to boiling with continuous stirring. If necessary, adjust the pH so that after sterilization it is about  $6.8 \pm 0.2$  at 25 °C. Sterilize in an autoclave using a validated cycle.

#### -Columbia agar

Pancreatic digest of casein 10.0 g

Meat peptic digest 5.0 g

Heart pancreatic digest 3.0 g

Yeast extract 5.0 g

Maize starch 1.0 g

Sodium chloride 5.0 g

Agar, according to gelling power 10.0 g to 15.0 g

Purified water 1000 mL

Hydrate the agar, dissolve by heating to boiling with continuous stirring. If necessary, adjust the pH so that after sterilization it is  $7.3 \pm 0.2$  at 25 °C. Sterilize in an autoclave using a validated cycle. Allow to cool to 45–50 °C; add, where necessary, gentamicin sulfate corresponding to 20 mg of gentamicin base and pour into Petri dishes.